

QUANTIFYING THE EFFECTS OF HIGHER ORDER JAHN-TELLER COUPLING TERMS ON A QUADRATIC JAHN-TELLER HAMILTONIAN IN THE CASE OF NO<sub>3</sub> AND Li<sub>3</sub>.

HENRY TRAN, *Department of Chemistry and Biochemistry, The Ohio State University, Columbus, OH, USA*; JOHN F. STANTON, *Department of Chemistry, The University of Texas, Austin, TX, USA*; TERRY A. MILLER, *Department of Chemistry and Biochemistry, The Ohio State University, Columbus, OH, USA*.

The Jahn-Teller (JT) effect represents an enormous complication in the understanding of many molecules. We have been able to assign  $\sim 20$  vibronic bands in the  $\tilde{A}^2E'' \leftarrow \tilde{X}^2A'_2$  transition of NO<sub>3</sub> and determine the linear and quadratic JT coupling terms for  $\nu_3$  and  $\nu_4$ , indicating strong and weak JT coupling along these modes respectively. It was found that the experimental results quantitatively disagree with ones determined from a vibronic Hamiltonian based on high-level *ab-initio* theory.<sup>a</sup> Typical analyses of experimental data use the quadratic JT Hamiltonian because limited measured levels tend to allow fitting only to coupling terms up to quadratic JT coupling. Hence, these analyses may neglect key contributions from cubic and quartic terms. To quantify this limitation, we have fit artificial spectra calculated with up to fourth order terms in the potential using a quadratic JT Hamiltonian and analyzed the results. The parameters chosen for this analysis are determined from *ab-initio* potentials for the  $\tilde{A}$  state of NO<sub>3</sub> and  $\tilde{X}$  state of Li<sub>3</sub> to gain further insight on these molecules. Our initial results concerning the limitations of the quadratic JT Hamiltonian will be presented.

<sup>a</sup>T. Codd, M.-W. Chen, M. Roudjane, J. F. Stanton, and T. A. Miller. Jet cooled cavity ringdown spectroscopy of the  $\tilde{A}^2E'' \leftarrow \tilde{X}^2A'_2$  Transition of the NO<sub>3</sub> Radical. *J. Chem. Phys.*, 142:184305, 2015